

140, Chemin de l'Armée d'Afrique
13010 Marseille
France
Tel : +33 (0) 4 96 12 20 40
Fax : +33 (0) 4 91 47 24 71
Email : info@biocytex.fr

DATA SHEET

Anti-GpVI PE

(Cat# 5131-PE100T)

MOUSE MONOCLONAL ANTIBODY ANTI HUMAN GpVI, R-PE CONJUGATED

Clone	1G5
Isotype	IgG2a, Kappa.
Immunogen	Recombinant human Glycoprotein VI (GpVI) ligand-binding ectodomain (residues 21-234).
Specificity	This antibody recognizes the human Glycoprotein VI (GpVI) ⁽¹⁾ . GpVI is a 62 kDa type 1 transmembrane receptor belonging to the immunoglobulin superfamily, and noncovalently associated with the signal-transducing FcRγ chain. GpVI is a major collagen receptor that plays a crucial role in the collagen-induced activation and aggregation of platelets. As well as collagen, other ligands such as collagen-related peptide (CRP) and snake venom protein (convulxin) can bind to the extracellular region of GpVI causing ectodomain shedding ⁽²⁾ . This process is tightly regulated by a metalloproteinase, most likely ADAM10, providing a mechanism for the modulation of platelet responsiveness. Most GpVI is maintained in a monomeric form on resting platelets. GpVI dimerization is a cAMP-controlled active process that primes platelet interaction with fibrillar collagen ⁽³⁾ .
Application	Flow Cytometry.
Form	R-Phycoerythrin conjugated purified immunoglobulin in PBS-BSA 0.1%, pH 7.2, liquid, 2 mL.
Size	100 tests, ready for use.
Suggested amount	20 µL/test per 10 ⁶ platelets in 100 µL sample.
Preservative	Sodium azide < 0.1%.
Storage	The conjugated antibody should be stored in the dark at +2-8°C until expiration date. Do not freeze.
References	<ol style="list-style-type: none">1. Al-Tamimi M. <i>et al.</i>, Platelets 2009, 20(2), 75-82.2. Gardiner E. <i>et al.</i>, J Thromb Haemost. 2007, 5,1530-1537.3. Loyau S. <i>et al.</i>, ATVB 2012, 32, 778-785.4. Qiao J. <i>et al.</i>, Blood 2013, 121:8, 1479-81.
Licence	Exclusive licence with Monash University.

FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR FOR DIAGNOSTIC PURPOSES.